arXiv:1407.1750v2 [math.RA] 18 Dec 2015

NON-ABELIAN TENSOR PRODUCT AND HOMOLOGY OF LIE SUPERALGEBRAS

X. GARCÍA-MARTÍNEZ, E. KHMALADZE, M. LADRA

Abstract. We introduce the non-abelian tensor product of Lie superalgebras and study some of its properties. We use it to describe the universal central extensions of Lie superalgebras. We present the low-dimensional non-abelian homology of Lie superalgebras and establish its relationship with the cyclic homology of associative superalgebras. We also define the non-abelian exterior product and give an analogue of Miller's theorem, Hopf formula and a six-term exact sequence for the homology of Lie superalgebras.

1. Introduction

In [1], Brown and Loday introduced the non-abelian tensor product of groups in the context of an application in homotopy theory. Analogous theories of non-abelian tensor product have been developed in other algebraic structures such as Lie algebras [8] and Lie–Rinehart algebras [4]. In [8], Ellis investigated the main properties of the non-abelian tensor product of Lie algebras and its relation to the low-dimensional homology of Lie algebras. In particular, he described the universal central extension of a perfect Lie algebra via the non-abelian tensor product. In [7], the non-abelian exterior product of Lie algebras is introduced and a six-term exact sequence relating low-dimensional homologies is obtained. In [10], using the non-abelian tensor product, Guin defined the non-abelian low-dimensional homology of Lie algebras and compared these groups with the cyclic homology and Milnor additive K-theory of associative algebras.

The theory of Lie superalgebras, also called $\mathbb{Z}_2$-graded Lie algebras, has aroused much interest both in mathematics and physics. Lie superalgebras play a very important role in theoretical physics since they are used to describe supersymmetry in a mathematical framework. A comprehensive description of the mathematical theory of Lie superalgebras is given in [14], containing the complete classification of all finite-dimensional simple Lie superalgebras over an algebraically closed field of characteristic zero. In the last few years, the theory of Lie superalgebras has experienced a remarkable evolution obtaining many results on representation theory and classification, most of them extending well-known facts on Lie algebras.

In this paper we develop the non-abelian tensor product and the low-dimensional non-abelian homology of Lie superalgebras, generalizing the corresponding notions for Lie algebras, with applications in universal central extensions and homology of Lie superalgebras and cyclic homology of associative superalgebras.

2010 *Mathematics Subject Classification.* 17B55, 17B30, 17B60.

Key words and phrases. Lie superalgebras, associative superalgebras, non-abelian tensor and exterior products, non-abelian homology, cyclic homology, Hopf formula, crossed module.

The organization of this paper is as follows: after this introduction, in Section 2 we give some definitions and necessary well-known results for the development of the paper. We also introduce actions and crossed modules of Lie superalgebras. In Section 3 we introduce the non-abelian tensor product of Lie superalgebras, we establish its principal properties such as right exactness and relation with the tensor product of supermodules. We describe the universal central extension of a perfect Lie superalgebra via the non-abelian tensor product (Theorem 4.1). In particular, applying this theorem, we obtain that $\mathfrak{st}(m,n,A)$ is the universal central extension of $\mathfrak{sl}(m,n,A)$, for $m+n \geq 5$, where A is a unital associative superalgebra. We also study nilpotency and solvability of the non-abelian tensor product of Lie superalgebras (Theorem 3.9). Using the non-abelian tensor product, in Section 5 we introduce the low-dimensional non-abelian homology of Lie superalgebras with coefficients in crossed modules. We show that, if the crossed module is a supermodule, then the non-abelian homology is the usual homology of Lie superalgebras. Then we apply this non-abelian homology to relate cyclic homology and Milnor cyclic homology of associative superalgebras, extending the results of [10]. Finally, in the last section we construct the non-abelian exterior product of Lie superalgebras and we use it to obtain Miller's type theorem for free Lie superalgebras, Hopf formula and a six-term exact sequence in the homology of Lie superalgebras.

Conventions and notations. Throughout this paper we denote by $\mathbb{K}$ a unital commutative ring unless otherwise stated. All modules and algebras are defined over $\mathbb{K}$. We write $\mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ and use its standard field structure. We put $(-1)^{\bar{0}} = 1$ and $(-1)^{\bar{1}} = -1$.

By *a supermodule* M we mean a module endowed with a $\mathbb{Z}_2$-gradation: $M = M_{\bar{0}} \oplus M_{\bar{1}}$. We call elements of $M_{\bar{0}}$ (resp. $M_{\bar{1}}$) even (resp. odd). Non-zero elements of $M_{\bar{0}} \cup M_{\bar{1}}$ will be called *homogeneous*. For a homogeneous $m \in M_{\bar{\alpha}}$, $\bar{\alpha} \in \mathbb{Z}_2$, its degree will be denoted by $|m|$. We adopt the convention that whenever the degree function occurs in a formula, the corresponding elements are supposed to be homogeneous. By *a homomorphism of supermodules* $f\colon M \to N$ of degree $|f| \in \mathbb{Z}_2$ we mean a linear map satisfying $f(M_{\bar{\alpha}}) \subseteq N_{\bar{\alpha}+|f|}$. In particular, if $|f| = \bar{0}$, then the homomorphism f will be called *of even grade (or even linear map)*.

By *a superalgebra* A we mean a supermodule $A = A_{\bar{0}} \oplus A_{\bar{1}}$ equipped with a bilinear multiplication satisfying $A_{\bar{\alpha}} A_{\bar{\beta}} \subseteq A_{\bar{\alpha}+\bar{\beta}}$, for $\bar{\alpha}, \bar{\beta} \in \mathbb{Z}_2$.

2. Preliminaries on Lie Superalgebras

In this section we review some terminology on Lie superalgebras and recall notions used in the paper. We mainly follow [2, 18], although with some modifications. We also introduce notions of actions and crossed modules of Lie superalgebras.

2.1. Definition and some examples of Lie superalgebras.

Definition 2.1. A *Lie superalgebra* is a superalgebra $M = M_{\bar{0}} \oplus M_{\bar{1}}$ with a multiplication denoted by $[\ ,\ ]$, called bracket operation, satisfying the following identities:

$$\begin{aligned} [x,y] &= -(-1)^{|x||y|}[y,x], \\ \big[x,[y,z]\big] &= \big[[x,y],z\big] + (-1)^{|x||y|}\big[y,[x,z]\big], \\ [m_{\bar{0}}, m_{\bar{0}}] &= 0, \end{aligned}$$

for all homogeneous elements $x, y, z \in M$ and $m_{\bar{0}} \in M_{\bar{0}}$.

Note that the last equation is an immediate consequence of the first one in the case 2 has an inverse in $\mathbb{K}$. Moreover, it can be easily seen that the second equation is equivalent to the graded Jacobi identity

$$(-1)^{|x||z|}\big[x,[y,z]\big]+(-1)^{|y||x|}\big[y,[z,x]\big]+(-1)^{|z||y|}\big[z,[x,y]\big]=0.$$

For a Lie superalgebra $M=M_{\bar{0}}\oplus M_{\bar{1}}$, the even part $M_{\bar{0}}$ is a Lie algebra. Hence, if $M_{\bar{1}}=0$, then M is just a Lie algebra. A Lie superalgebra M without even part, i. e., $M_{\bar{0}}=0$, is an *abelian Lie superalgebra*, that is, $[x,y]=0$ for all $x,y\in M$.

A *Lie superalgebra homomorphism* $f\colon M\to M'$ is a supermodule homomorphism of even grade such that $f[x,y]=[f(x),f(y)]$ for all $x,y\in M$.

Example 2.2.

(i) Any associative superalgebra A can be considered as a Lie superalgebra with the bracket

$$[a,b]=ab-(-1)^{|a||b|}ba.$$

(ii) Let m, n be positive integers and A a unital associative superalgebra. Consider the algebra $\mathcal{M}(m,n,A)$ of all $(m+n)\times(m+n)$-matrices with entries in A and with the usual product of matrices. A $\mathbb{Z}_2$-gradation is defined as follows: homogeneous elements are matrices $E_{ij}(a)$ having the homogeneous element $a\in A$ at the position (i,j) and zero elsewhere, and $|E_{ij}(a)|=|i|+|j|+|a|$, where $|i|=\bar{0}$ if $1\le i\le m$ and $|i|=\bar{1}$ if $m+1\le i\le m+n$. With this gradation, $\mathcal{M}(m,n,A)$ turns out to be an associative superalgebra. The corresponding Lie superalgebra will be denoted by $\mathfrak{gl}(m,n,A)$.

(iii) Let $V=V_{\bar{0}}\oplus V_{\bar{1}}$ be a supermodule. Then the supermodule $\mathrm{End}_{\mathbb{K}}(V)$ of all linear endomorphisms $V\to V$ (of both degrees 0 and 1) has a structure of an associative superalgebra with respect to composition (see [2]) and hence becomes a Lie superalgebra. In particular, if the ground ring $\mathbb{K}$ is a field, and m, n are dimensions of $V_{\bar{0}}$ and $V_{\bar{1}}$ respectively, then choosing a homogeneous basis of V ordered such that even elements stand before odd, the elements of $\mathrm{End}_{\mathbb{K}}(V)$ can be seen as $(m+n)\times(m+n)$-square matrices

$$\begin{pmatrix} a & b\\ c & d\end{pmatrix}$$

where a,b,c and d are respectively $m\times m$, $m\times n$, $n\times m$ and $n\times n$ matrices with entries in $\mathbb{K}$. The even elements are the matrices with $b=c=0$ and the odd elements are matrices with $a=d=0$.

Let M and N be two submodules of a Lie superalgebra P. We denote by $[M,N]$ the submodule of P spanned by all elements $[m,n]$ with $m\in M$ and $n\in N$. A $\mathbb{Z}_2$-graded submodule M is a *graded ideal* of P if $[M,P]\subseteq M$. In particular, the submodule $\mathsf{Z}(P)=\{c\in P:[c,p]=0 \text{ for all } p\in P\}$ is a graded ideal and it is called the *centre* of P. Clearly if M and N are graded ideals of P, then so is $[M,N]$.

Let M be a Lie superalgebra and $D\in\mathrm{End}_{\mathbb{K}}(M)$. We say that D is a *derivation* if for all $x,y\in M$

$$D([x,y])=[D(x),y]+(-1)^{|D||x|}[x,D(y)].$$

We denote by $\big(\mathrm{Der}_{\mathbb{K}}(M)\big)_{\bar{\alpha}}$ the set of homogeneous derivations of degree $\bar{\alpha}\in\mathbb{Z}_2$. One verifies that the supermodule of derivations

$$\mathrm{Der}_{\mathbb{K}}(M)=\big(\mathrm{Der}_{\mathbb{K}}(M)\big)_{\bar{0}}\oplus\big(\mathrm{Der}_{\mathbb{K}}(M)\big)_{\bar{1}}$$

is a subalgebra of the Lie superalgebra $\mathrm{End}_{\mathbb{K}}(M)$.

2.2. **Actions and crossed modules of Lie superalgebras.**

Definition 2.3. Let P and M be two Lie superalgebras. By an *action* of P on M we mean a $\mathbb{K}$-bilinear map of even grade,

$$P \times M \to M, \quad (p, m) \mapsto {}^{p}m,$$

such that

(i) ${}^{[p,p']}m = {}^{p}({}^{p'}m) - (-1)^{|p||p'|}\, {}^{p'}({}^{p}m)$,
(ii) ${}^{p}[m, m'] = [{}^{p}m, m'] + (-1)^{|p||m|}[m, {}^{p}m']$,

for all homogeneous $p, p' \in P$ and $m, m' \in M$.

The action is called *trivial* if ${}^{p}m = 0$ for all $p \in P$ and $m \in M$.

For example, if M is a graded ideal and P is a subalgebra of a Lie superalgebra Q, then the bracket in Q induces an action of P on M.

Note that the action of P on M is the same as a Lie superalgebra homomorphism $P \to \mathrm{Der}_{\mathbb{K}}(M)$.

Remark. If M is an abelian Lie superalgebra enriched with an action of a Lie superalgebra P, then M has a structure of a supermodule over P (P-supermodule, for short) (see e. g. [18]), that is, there is a $\mathbb{K}$-bilinear map of even grade $P \times M \to M$, $(p, m) \mapsto pm$, such that

$$[p, p']m = p(p'm) - (-1)^{|p||p'|}p'(pm),$$

for all homogeneous $p, p' \in P$ and $m \in M$.

Note that a P-supermodule M is the same as a $\mathbb{K}$-supermodule M together with a Lie superalgebra homomorphism $P \to \mathrm{End}_{\mathbb{K}}(M)$.

Definition 2.4. Given two Lie superalgebras M and P with an action of P on M, we can define the *semidirect product* $M \rtimes P$ with the underlying supermodule $M \oplus P$ endowed with the bracket given by

$$[(m, p), (m', p')] = ([m, m'] + {}^{p}m' - (-1)^{|m||p'|}({}^{p'}m), [p, p']).$$

Now we are ready to introduce the following notion of crossed modules of Lie superalgebras (see also [23, Definition 5]).

Definition 2.5. A *crossed module of Lie superalgebras* is a homomorphism of Lie superalgebras $\partial\colon M \to P$ with an action of P on M satisfying

(i) $\partial({}^{p}m) = [p, \partial(m)]$,
(ii) ${}^{\partial(m)}m' = [m, m']$,

for all $p \in P$ and $m, m' \in M$.

Example 2.6. There are some standard examples of crossed modules:

(i) The inclusion $M \hookrightarrow P$ of a graded ideal M of a Lie superalgebra P is a crossed module of Lie superalgebras.

(ii) If P is a Lie superalgebra and M is a P-supermodule, the trivial map $0\colon M \to P$ is a crossed module of Lie superalgebras.

(iii) A central extension of Lie superalgebras $\partial\colon M \twoheadrightarrow P$ (i.e., $\mathsf{Ker}\,\partial \subseteq \mathsf{Z}(M)$) is a crossed module of Lie superalgebras. Here the action of P on M is given by ${}^{p}m = [\bar{m}, m]$, where $\bar{m} \in M$ is any element of $\partial^{-1}(p)$.

(iv) The homomorphism of Lie superalgebras $\partial\colon M \to \mathrm{Der}_{\mathbb{K}}(M)$ which sends $m \in M$ to the inner derivation $\mathsf{ad}(m) \in \mathrm{Der}_{\mathbb{K}}(M)$, defined by $\mathsf{ad}(m)(m') = [m, m']$, together with the action of $\mathrm{Der}_{\mathbb{K}}(M)$ on M given by ${}^{D}m = D(m)$, is a crossed module of Lie superalgebras.

Lemma 2.7. *Let $\partial\colon M \to P$ be a crossed module of Lie superalgebras. Then the following conditions are satisfied:*

(i) *The kernel of ∂ is in the centre of M.*
(ii) *The image of ∂ is a graded ideal of P.*
(iii) *The Lie superalgebra $\mathsf{Im}\,\partial$ acts trivially on the centre $\mathsf{Z}(M)$, and so trivially on $\mathsf{Ker}\,\partial$. Hence $\mathsf{Ker}\,\partial$ inherits an action of $P/\mathsf{Im}\,\partial$ making $\mathsf{Ker}\,\partial$ a $P/\mathsf{Im}\,\partial$-supermodule.*

Proof. This is an immediate consequence of Definition 2.5. □

2.3. **Free Lie superalgebra and enveloping superalgebra of a Lie superalgebra.**

Definition 2.8. The *free Lie superalgebra* on a $\mathbb{Z}_2$-graded set $X = X_{\bar{0}} \cup X_{\bar{1}}$ is a Lie superalgebra $\mathrm{F}(X)$ together with a degree zero map $i\colon X \to \mathrm{F}(X)$ such that if M is any Lie superalgebra and $j\colon X \to M$ is a degree zero map, then there is a unique Lie superalgebra homomorphism $h\colon \mathrm{F}(X) \to M$ with $j = h \circ i$.

The existence of free Lie superalgebras is guaranteed by an analogue of Witt's theorem (see [18, Theorem 6.2.1]). In the sequel we need the following construction of the free Lie superalgebra.

Construction 2.9. *Let $X = X_{\bar{0}} \cup X_{\bar{1}}$ be a $\mathbb{Z}_2$-graded set. Denote by $\mathrm{mag}(X)$ the free magma over the set X. The free superalgebra on X, denoted by $\mathrm{alg}(X)$, has as elements the finite sums $\sum_i \lambda_i v_i$, where $\lambda_i \in \mathbb{K}$ and x_i are elements of $\mathrm{mag}(X)$ and the multiplication in $\mathrm{alg}(X)$ extends the multiplication in $\mathrm{mag}(X)$. Note that the grading is naturally defined in $\mathrm{alg}(X)$. The free Lie superalgebra $\mathrm{F}(X)$ is the quotient $\mathrm{alg}(X)/I$, where I is the graded ideal generated by the elements*

$$xy + (-1)^{|x||y|}yx,$$
$$(-1)^{|x||z|}\big(x(yz)\big) + (-1)^{|y||x|}\big(y(zx)\big) + (-1)^{|z||y|}\big(z(xy)\big),$$
$$x_{\bar{0}}x_{\bar{0}},$$

for all homogeneous $x, y, z \in X$ and $x_{\bar{0}} \in X_{\bar{0}}$.

Definition 2.10. The *universal enveloping superalgebra* of a Lie superalgebra M is a pair $(\mathrm{U}(M), \sigma)$, where $\mathrm{U}(M)$ is a unital associative superalgebra and $\sigma\colon M \to \mathrm{U}(M)$ is an even linear map satisfying

$$\sigma[x, y] = \sigma(x)\sigma(y) - (-1)^{|x||y|}\sigma(y)\sigma(x), \tag{1}$$

for all homogeneous $x, y \in M$, such that the following universal property holds: for any other pair (A, σ'), where A is a unital associative superalgebra and $\sigma'\colon M \to A$ is an even linear map satisfying (1), there is a unique superalgebra homomorphism $f\colon \mathrm{U}(M) \to A$ such that $f \circ \sigma = \sigma'$.

Now we need to recall (see e. g. [21]) that, given two supermodules M and N, the tensor product of modules $M \otimes_{\mathbb{K}} N$ has a natural supermodule structure with

$\mathbb{Z}_2$-grading given by

$$(M \otimes_{\mathbb{K}} N)_{\bar{\alpha}} = \bigoplus_{\bar{\beta}+\bar{\gamma}=\bar{\alpha}} (M_{\bar{\beta}} \otimes_{\mathbb{K}} N_{\bar{\gamma}}).$$

In particular, the tensor power $M^{\otimes n}$, $n \geq 2$, has the induced $\mathbb{Z}_2$-grading. Hence the tensor algebra $T(M)$ has the $\mathbb{Z}_2$-grading extending that of M. We call $T(M)$ *the tensor superalgebra*.

Construction 2.11. *Let M be a Lie superalgebra and $T(M)$ the tensor superalgebra over the underlying supermodule of M. Consider the two-sided ideal $J(M)$ of $T(M)$ generated by all elements of the form*

$$m \otimes m' - (-1)^{|m||m'|} m' \otimes m - [m, m'],$$

for all homogeneous $m, m' \in M$. Then the quotient $\mathrm{U}(M) = T(M)/J(M)$ is a unital associative superalgebra. By composing the canonical inclusion $M \to T(M)$ with the canonical projection $T(M) \to \mathrm{U}(M)$ we get the canonical even linear map $\sigma\colon M \to \mathrm{U}(M)$. Then the pair $(\mathrm{U}(M), \sigma)$ is the universal enveloping superalgebra of M (see [2]*).*

Note that, as in the Lie algebra case, the universal enveloping superalgebra turns out to be a very useful tool for the representation theory of Lie superalgebras. In particular, by the universal property, it follows that a Lie supermodule over a Lie superalgebra M is the same as a $\mathbb{Z}_2$-graded (left) $\mathrm{U}(M)$-module (see [21, Chapter 1]).

Let us consider $\mathbb{K}$ with $\mathbb{Z}_2$-grading concentrated in degree zero, that is, with $\mathbb{K}_{\bar{1}} = 0$. Then the trivial map from a Lie superalgebra M into $\mathbb{K}$ gives rise to a unique homomorphism of superalgebras $\varepsilon\colon \mathrm{U}(M) \to \mathbb{K}$. The kernel of ε, denoted by $\Omega(M)$, is called the *augmentation ideal* of M. Obviously, $\Omega(M)$ is just the graded ideal of $\mathrm{U}(M)$ generated by $\sigma(M)$.

2.4. **Homology of Lie superalgebras.** Now we briefly recall from [18, 22] the definition of homology of Lie superalgebras.

The *Grassmann algebra* of a Lie superalgebra P, denoted by $\bigwedge_{\mathbb{K}}(P)$, is defined to be the quotient of the tensor superalgebra $T(P)$ of P by the ideal generated by the elements

$$x \otimes y + (-1)^{|x||y|} y \otimes x,$$

for all homogeneous $x, y \in P$. Note that $\bigwedge_{\mathbb{K}}(P) = \bigoplus_{n>0} \bigwedge^n_{\mathbb{K}}(P)$, where $\bigwedge^n_{\mathbb{K}}(P)$ is the image of $P^{\otimes n}$ in $\bigwedge_{\mathbb{K}}(P)$, has an induced P-supermodule structure given by

$$x(x_1 \wedge \cdots \wedge x_n) = \sum_{i=1}^{n} (-1)^{|x| \sum_{k<i} |x_k|} (x_1 \wedge \cdots \wedge [x, x_i] \wedge \cdots \wedge x_n).$$

Let M be a P-supermodule and consider the chain complex $(C_*(P, M), d_*)$ defined by $C_n(P, M) = \bigwedge^n_{\mathbb{K}}(P) \otimes_{\mathbb{K}} M$, for $n \geq 0$, with boundary maps $d_n\colon C_n(P, M) \to C_{n-1}(P, M)$ defined on generators by

$$\begin{aligned} d_n(x_1 \wedge \cdots \wedge x_n \otimes y) &= \sum_{i=1}^{n} (-1)^{i+|x_i| \sum_{k>i} |x_k|} (x_1 \wedge \cdots \wedge \hat{x}_i \wedge \cdots \wedge x_n \otimes x_i y) \\ &+ \sum_{i<j} (-1)^{i+j+|x_i| \sum_{k<i} |x_k| + |x_j| \sum_{l<j} |x_l| + |x_i||x_j|} ([x_i, x_j] \wedge \cdots \wedge \hat{x}_i \wedge \cdots \wedge \hat{x}_j \wedge \cdots \wedge x_n \otimes y). \end{aligned}$$

The n-th *homology of the Lie superalgebra* P with coefficients in the P-supermodule M, $\mathrm{H}_n(P,M)$, is the n-th homology of the chain complex $(C_*(P,M),d_*)$, i.e.

$$\mathrm{H}_n(P,M)=\frac{\mathsf{Ker}\,d_n}{\mathsf{Im}\,d_{n+1}}.$$

If $\mathbb{K}$ is regarded as a trivial P-supermodule, we write $\mathrm{H}_n(P)$ for $\mathrm{H}_n(P,\mathbb{K})$.

In the case when the ground ring $\mathbb{K}$ is a field, there is a relation between Tor functor and the homology (see [18]) given by

$$\mathrm{H}_n(P,M)\cong\mathsf{Tor}_n^{\mathrm{U}(P)}(\mathbb{K},M).$$

By analogy to Lie algebras (see e. g. [11]), we have the following isomorphisms

$$H_0(P,M)\cong\mathsf{Coker}\big(\Omega(P)\otimes_{\mathrm{U}(P)}M\longrightarrow M\big), \tag{2}$$

$$H_1(P,M)\cong\mathsf{Ker}\big(\Omega(P)\otimes_{\mathrm{U}(P)}M\longrightarrow M\big). \tag{3}$$

3. Non-abelian tensor product of Lie superalgebras

In this section we introduce a non-abelian tensor product of Lie superalgebras, which generalizes the non-abelian tensor product of Lie algebras [8], and study its properties.

3.1. **Construction of the non-abelian tensor product.**

Definition 3.1. Let M and N be two Lie superalgebras with actions on each other. Let $X_{M,N}$ be the $\mathbb{Z}_2$-graded set of all symbols $m\otimes n$, where $m\in M_{\bar0}\cup M_{\bar1}$, $n\in N_{\bar0}\cup N_{\bar1}$ and the $\mathbb{Z}_2$-gradation is given by $|m\otimes n|=|m|+|n|$. We define the *non-abelian tensor product* of M and N, denoted by $M\otimes N$, as the Lie superalgebra generated by $X_{M,N}$ and subject to the relations:

(i) $\lambda(m\otimes n)=\lambda m\otimes n=m\otimes\lambda n$,

(ii) $(m+m')\otimes n=m\otimes n+m'\otimes n$, where m,m' have the same grade,

$m\otimes(n+n')=m\otimes n+m\otimes n'$, where n,n' have the same grade,

(iii) $[m,m']\otimes n=m\otimes{}^{m'}n-(-1)^{|m||m'|}(m'\otimes{}^{m}n)$,

$m\otimes[n,n']=(-1)^{|n'|(|m|+|n|)}({}^{n'}m\otimes n)-(-1)^{|m||n|}({}^{n}m\otimes n')$,

(iv) $[m\otimes n,m'\otimes n']=-(-1)^{|m||n|}({}^{n}m\otimes{}^{m'}n')$,

for every $\lambda\in\mathbb{K}$, $m,m'\in M_{\bar0}\cup M_{\bar1}$ and $n,n'\in N_{\bar0}\cup N_{\bar1}$.

Let us remark that if $m=m_{\bar0}+m_{\bar1}$ is any element of M and $n=n_{\bar0}+n_{\bar1}$ is any element of N, then under the notation $m\otimes n$ we mean the sum

$$m_{\bar0}\otimes n_{\bar0}+m_{\bar0}\otimes n_{\bar1}+m_{\bar1}\otimes n_{\bar0}+m_{\bar1}\otimes n_{\bar1}.$$

If $M=M_{\bar0}$ and $N=N_{\bar0}$ then $M\otimes N$ is the non-abelian tensor product of Lie algebras introduced and studied in [8] (see also [12]).

Definition 3.2. Actions of Lie superalgebras M and N on each other are said to be *compatible* if

(i) ${}^{({}^{n}m)}n'=-(-1)^{|m||n|}[{}^{m}n,n']$,

(ii) ${}^{({}^{m}n)}m'=-(-1)^{|m||n|}[{}^{n}m,m']$,

for all $m,m'\in M_{\bar0}\cup M_{\bar1}$ and $n,n'\in N_{\bar0}\cup N_{\bar1}$.

For example, if M and N are two graded ideals of some Lie superalgebra, the actions induced by the bracket are compatible.

Proposition 3.3. *Let M and N be Lie superalgebras acting compatibly on each other. Then there is a natural isomorphism of Lie superalgebras*

$$M \otimes N \cong \frac{M \otimes_{\mathbb{K}} N}{D(M,N)},$$

where $D(M,N)$ is the submodule of the supermodule $M \otimes_{\mathbb{K}} N$ generated by the elements

(i) $[m,m'] \otimes n - m \otimes {}^{m'}n + (-1)^{|m||m'|}(m' \otimes {}^{m}n)$,
(ii) $m \otimes [n,n'] - (-1)^{|n'|(|m|+|n|)}({}^{n'}m \otimes n) + (-1)^{|m||n|}({}^{n}m \otimes n')$,
(iii) $({}^{n}m) \otimes ({}^{m}n)$, *with* $|m| = |n|$
(iv) $(-1)^{|m||n|}({}^{n}m) \otimes ({}^{m'}n') + (-1)^{(|m|+|n|)(|m'|+|n'|)+|m'||n'|}({}^{n'}m') \otimes ({}^{m}n)$,
(v) $\displaystyle\bigcirclearrowleft_{(m,n),(m',n'),(m'',n'')} (-1)^{(|m|+|n|)(|m''|+|n''|)+|m||n|+|m'||n'|}[{}^{n}m, {}^{n'}m'] \otimes ({}^{m''}n'')$,

for all $m,m',m'' \in M_{\bar{0}} \cup M_{\bar{1}}$ and $n,n',n'' \in N_{\bar{0}} \cup N_{\bar{1}}$, where $\displaystyle\bigcirclearrowleft_{x,y,z}$ denotes the cyclic summation with respect to x,y,z.

Proof. There is a Lie superalgebra structure on the supermodule $(M \otimes_{\mathbb{K}} N)/D(M,N)$ given on generators by the following bracket

$$[m \otimes n, m' \otimes n'] = -(-1)^{|m||n|}({}^{n}m \otimes {}^{m'}n'),$$

for all $m,m' \in M_{\bar{0}} \cup M_{\bar{1}}$, $n,n' \in N_{\bar{0}} \cup N_{\bar{1}}$ and extended by linearity. It is routine to check that this bracket is compatible with the defining relations of $(M \otimes_{\mathbb{K}} N)/D(M,N)$ and it indeed defines a Lie superalgebra structure. Then the canonical homomorphism $M \otimes N \to (M \otimes_{\mathbb{K}} N)/D(M,N)$, $m \otimes n \mapsto m \otimes n$, is an isomorphism. □

The proof of the following proposition is a routine calculation.

Proposition 3.4. *Let M and N be two Lie superalgebras acting compatibly on each other.*

(i) *The following morphisms*

$$\mu\colon M \otimes N \to M, \quad m \otimes n \mapsto -(-1)^{|m||n|}({}^{n}m),$$
$$\nu\colon M \otimes N \to N, \quad m \otimes n \mapsto {}^{m}n,$$

are Lie superalgebra homomorphisms.

(ii) *There are actions of M and N on $M \otimes N$ given by*

$${}^{m'}(m \otimes n) = [m',m] \otimes n + (-1)^{|m||m'|}m \otimes ({}^{m'}n),$$
$${}^{n'}(m \otimes n) = ({}^{n'}m) \otimes n + (-1)^{|n||n'|}m \otimes [n',n],$$

for $m,m' \in M_{\bar{0}} \cup M_{\bar{1}}$, $n,n' \in N_{\bar{0}} \cup N_{\bar{1}}$ and extended by linearity. Moreover, with these actions μ and ν are crossed modules of Lie superalgebras.

We will denote by $[M,N]^M$ (resp. $[M,N]^N$) the image of μ (resp. ν), which by Lemma 2.7(ii) is a graded ideal of M (resp. N) generated by the elements of the form ${}^{n}m$ (resp. ${}^{m}n$) for $m \in M$ and $n \in N$. Note that by Lemma 2.7(iii) $\mathsf{Ker}(\mu)$ $\big($resp. $\mathsf{Ker}(\nu)\big)$ is an $M/[M,N]^M$-supermodule (resp. $N/[M,N]^N$-supermodule).

3.2. **Some properties of the non-abelian tensor product.** The obvious analogues of Brown and Loday results [1] hold for Lie superalgebras. In the following two propositions immediately below we show that sometimes the non-abelian tensor product of Lie superalgebras can be expressed in terms of the tensor product of supermodules.

Proposition 3.5. *Let M and N be Lie superalgebras acting on each other. Then the canonical map $M \otimes_{\mathbb{K}} N \to M \otimes N$, $m \otimes n \mapsto m \otimes n$, is an even, surjective homomorphism of supermodules. In addition, if M and N act trivially on each other, then $M \otimes N$ is an abelian Lie superalgebra and there is an isomorphism of supermodules*

$$M \otimes N \cong M^{\mathrm{ab}} \otimes_{\mathbb{K}} N^{\mathrm{ab}},$$

where $M^{\mathrm{ab}} = M/[M,M]$ and $N^{\mathrm{ab}} = N/[N,N]$.

Proof. It is straightforward by the identities (iv), (iii) of Definition 3.1. □

Proposition 3.6. *Let P be a Lie superalgebra and M a P-supermodule considered as an abelian Lie superalgebra acting trivially on P. Then there is an isomorphism of supermodules*

$$P \otimes M \cong \Omega(P) \otimes_{\mathrm{U}(P)} M.$$

Proof. By Proposition 3.3 there is an isomorphism of supermodules

$$P \otimes M \cong \frac{P \otimes_{\mathbb{K}} M}{W},$$

where W is the submodule of $P \otimes_{\mathbb{K}} M$ generated by all elements of the form

$$[p,p'] \otimes m - p \otimes p'm + (-1)^{|p||p'|} p' \otimes pm$$

for all $p, p' \in P_{\bar{0}} \cup P_{\bar{1}}$ and $m \in M_{\bar{0}} \cup M_{\bar{1}}$. Now by using Construction 2.11 and by repeating the respective part of the proof of [3, Proposition 13], it is easy to see that there is an isomorphism of supermodules

$$\frac{P \otimes_{\mathbb{K}} M}{W} \cong \Omega(P) \otimes_{\mathrm{U}(P)} M,$$

which completes the proof. □

The non-abelian tensor product of Lie superalgebras is symmetric, in the sense of the following proposition.

Proposition 3.7. *The Lie superalgebra homomorphism*

$$M \otimes N \to N \otimes M, \quad m \otimes n \mapsto -(-1)^{|m||n|}(n \otimes m),$$

is an isomorphism.

Proof. This can be checked readily. □

Let us consider the category $\mathtt{SLie}^2_{\mathbb{K}}$ whose objects are ordered pairs of Lie superalgebras (M,N) acting compatibly on each other, and the morphisms are pairs of Lie superalgebra homomorphisms $(\phi\colon M \to M', \psi\colon N \to N')$ which preserve the actions, i.e., $\phi({}^n m) = {}^{\psi(n)}\phi(m)$ and $\psi({}^m n) = {}^{\phi(m)}\psi(n)$. For such a pair (ϕ,ψ) we have a homomorphism of Lie superalgebras $\phi \otimes \psi\colon M \otimes N \to M' \otimes N'$, $m \otimes n \mapsto \phi(m) \otimes \psi(n)$. Therefore, $\otimes$ is a functor from $\mathtt{SLie}^2_{\mathbb{K}}$ to the category of Lie superalgebras.

Given an exact sequence in $\mathtt{SLie}^2_{\mathbb{K}}$

$$(0,0) \longrightarrow (K,L) \xrightarrow{(i,j)} (M,N) \xrightarrow{(\phi,\psi)} (P,Q) \longrightarrow (0,0), \tag{4}$$

by Proposition 3.4(ii) there is a Lie superalgebra homomorphism $M \otimes L \to L$ and an action of N on $K \otimes N$. Thus, there is an action of $M \otimes L$ on $K \otimes N$, so we can form the semidirect product $(K \otimes N) \rtimes (M \otimes L)$, and we have the following obvious analogue of [8, Proposition 9].

Proposition 3.8. *Given the short exact sequence* (4)*, there is an exact sequence of Lie superalgebras*

$$(K \otimes N) \rtimes (M \otimes L) \xrightarrow{\alpha} M \otimes N \xrightarrow{\phi\otimes\psi} P \otimes Q \longrightarrow 0.$$

In particular, given a Lie superalgebra M and a graded ideal K of M, there is an exact sequences of Lie superalgebras

$$(K \otimes M) \rtimes (M \otimes K) \to M \otimes M \to (M/K) \otimes (M/K) \to 0. \tag{5}$$

3.3. **Nilpotency, solvability and Engel of the non-abelian tensor product.** The results from [20] on nilpotency, solvability and Engel of the non-abelian tensor product on Lie algebras can be easily extended to the case of Lie superalgebras. The notions of nilpotency and solvability of Lie superalgebras are given in [18]. As they are very similar to the respective notions for Lie algebras, we omit them. We say that a Lie superalgebra M is *n-Engel* if it satisfies $\mathsf{ad}(x)^n = 0$ for all $x \in M$. The proof of the following result is similar to the proof of [20, Theorem 2.2].

Theorem 3.9. *Let M and N be two Lie superalgebras acting compatibly on each other. Then,*

(i) *If $[M,N]^M$ is nilpotent, then $M \otimes N$ and $[M,N]^N$ are nilpotent too. Moreover, if the nilpotency class of $[M,N]^M$ is $\mathrm{cl}([M,N]^M)$, then*

$$\mathrm{cl}([M,N]^M) \le \mathrm{cl}(M \otimes N) \le \mathrm{cl}([M,N]^M) + 1,$$
$$\mathrm{cl}([M,N]^N) \le \mathrm{cl}([M,N]^M) + 1.$$

(ii) *If $[M,N]^M$ is solvable, then $M \otimes N$ and $[M,N]^N$ are solvable too. Moreover, if the derived length of $[M,N]^M$ is $\ell([M,N]^M)$, then*

$$\ell([M,N]^M) \le \ell(M \otimes N) \le \ell([M,N]^M) + 1,$$
$$\ell([M,N]^N) \le \ell([M,N]^M) + 1.$$

(iii) *If $[M,N]^M$ is Engel, then $M \otimes N$ and $[M,N]^N$ are Engel too. Moreover, if $[M,N]^M$ is n-Engel, then $M \otimes N$ and $[M,N]^N$ are $(n+1)$-Engel.*

4. Universal central extensions of Lie superalgebras

Now we use the non-abelian tensor product of Lie superalgebras to describe universal central extensions of Lie superalgebras. Recall that a central extension $\mathfrak{u}\colon U \twoheadrightarrow P$ is universal if for any other central extension $f\colon M \twoheadrightarrow P$ there is a unique homomorphism $\theta\colon U \to M$ such that $f \circ \theta = \mathfrak{u}$. It is shown in [19] that a Lie superalgebra P admits a universal central extension if and only if P is perfect, i.e. $P = [P,P]$.

It follows from Proposition 3.4 and Lemma 2.7(i) that the homomorphism $\mathfrak{u}\colon P \otimes P \twoheadrightarrow [P,P]$, $\mathfrak{u}(p \otimes p') = [p,p']$, is a central extension of the Lie superalgebra $[P,P]$.

Theorem 4.1. *If P is a perfect Lie superalgebra, then the central extension $\mathfrak{u}\colon P\otimes P\twoheadrightarrow P$ is the universal central extension.*

Proof. Let $f\colon M\twoheadrightarrow P$ be a central extension of P. Since $\mathsf{Ker}\, f$ is in the centre of M, we get a well-defined homomorphism of Lie superalgebras $\theta\colon P\otimes P\to M$ given by $\theta(p\otimes p')=[m_p,m_{p'}]$, where m_p and $m_{p'}$ are any preimages of p and p', respectively. Obviously $\theta\circ f=\mathfrak{u}$. Since P is perfect, then by relation (iv) of Definition 3.1, so is $P\otimes P$. Then by [19, Lemma 1.4] the homomorphism θ is unique. □

Remark. If P is a perfect Lie superalgebra, then $\mathrm{H}_2(P)\approx\mathsf{Ker}(P\otimes P\xrightarrow{\mathfrak{u}}P)$, since the kernel of the universal central extension is isomorphic to the second homology $\mathrm{H}_2(P)$ (see [19]).

It is a classical result that the universal central extension of the Lie algebra $\mathfrak{sl}(n,A)$, where A is a unital associative algebra, is the Steinberg algebra $\mathfrak{st}(n,A)$, when $n\geq 5$ (see e. g. [16]). Recently, in [5, 9], this result has been extended to Lie superalgebras. Below, using the non-abelian tensor product of Lie superalgebras, we propose an alternative proof of the same result.

First we recall from [5] that, given a unital associative superalgebra A, the *Lie superalgebra* $\mathfrak{sl}(m,n,A)$, $m+n\geq 3$, is defined to be the subalgebra of the Lie superalgebra $\mathfrak{gl}(m,n,A)$ (see Example 2.2 (ii)) generated by the elements $E_{ij}(a)$, $1\leq i\neq j\leq m+n$, $a\in A_{\bar{0}}\cup A_{\bar{1}}$. It is shown in [5, Lemma 3.3] that $\mathfrak{sl}(m,n,A)$ is a perfect Lie superalgebra. This guarantees the existence of the universal central extension of $\mathfrak{sl}(m,n,A)$.

The Steinberg Lie superalgebra $\mathfrak{st}(m,n,A)$ is defined for $m+n\geq 3$ to be the Lie superalgebra generated by the homogeneous elements $F_{ij}(a)$, where $1\leq i\neq j\leq m+n$, $a\in A$ is a homogeneous element and the $\mathbb{Z}_2$-grading is given by $|F_{ij}(a)|=|i|+|j|+|a|$, subject to the following relations:

$$\begin{aligned}
&a\mapsto F_{ij}(a) \text{ is a } \mathbb{K}\text{-linear map},\\
&[F_{ij}(a),F_{jk}(b)]=F_{ik}(ab), \text{ for distinct } i,j,k,\\
&[F_{ij}(a),F_{kl}(b)]=0, \text{ for } j\neq k, i\neq l.
\end{aligned}$$

Theorem 4.2 ([5])**.** *If $m+n\geq 5$, then the canonical epimorphism*

$$\mathfrak{st}(m,n,A)\twoheadrightarrow\mathfrak{sl}(m,n,A),\quad F_{ij}(a)\mapsto E_{ij}(a),$$

is the universal central extension of the perfect Lie superalgebra $\mathfrak{sl}(m,n,A)$.

Proof. We claim that there is an isomorphism of Lie superalgebras

$$\mathfrak{st}(m,n,A)\cong\mathfrak{st}(m,n,A)\otimes\mathfrak{st}(m,n,A).$$

Indeed, one can readily check that the maps

$$\begin{aligned}
&\mathfrak{st}(m,n,A)\longrightarrow\mathfrak{st}(m,n,A)\otimes\mathfrak{st}(m,n,A),\quad F_{ij}(a)\mapsto F_{ik}(a)\otimes F_{kj}(1) \text{ for } k\neq i,j,\\
&\mathfrak{st}(m,n,A)\otimes\mathfrak{st}(m,n,A)\longrightarrow\mathfrak{st}(m,n,A),\quad F_{ij}(a)\otimes F_{kl}(b)\mapsto[F_{ij}(a),F_{kl}(b)],
\end{aligned}$$

are well-defined homomorphisms of Lie superalgebras if $m+n\geq 5$, and they are inverses to each other. Since $\mathfrak{st}(m,n,A)$ is a perfect Lie superalgebra, then Theorem 4.1 and [19, Corollary 1.9] complete the proof. □

5. Non-abelian homology of Lie superalgebras

The low-dimensional non-abelian homology of Lie algebras with coefficients in crossed modules was defined in [10] and it was extended to all dimensions in [12]. In this section we extend to Lie superalgebras the construction of zero and first non-abelian homologies. We also relate the non-abelian homology of Lie superalgebras with the cyclic homology of associative superalgebras studied in [13, 15].

5.1. **Construction of the non-abelian homology and some properties.** Let P be a Lie superalgebra. We denote by $\mathtt{Cross}(\mathrm{P})$ the category of crossed modules of Lie superalgebras over P (crossed P-modules, for short), whose objects are crossed modules $(M,\partial) \equiv (\partial\colon M \to P)$ and a morphism from (M,∂) to (N,∂') is a Lie superalgebra homomorphism $f\colon M \to N$ such that $f({}^p m) = {}^p f(m)$ for all $p \in P$, $m \in M$ and $\partial' \circ f = \partial$. By an exact sequence $(L,\partial'') \xrightarrow{f} (M,\partial) \xrightarrow{g} (N,\partial')$ in $\mathtt{Cross}(\mathrm{P})$ we mean that the sequence of Lie superalgebras $L \xrightarrow{f} M \xrightarrow{g} N$ is exact.

Lemma 5.1. *Given a short exact sequence in* $\mathtt{Cross}(\mathrm{P})$

$$0 \to (L,\partial'') \xrightarrow{f} (M,\partial) \xrightarrow{g} (N,\partial') \to 0,$$

the morphism $\partial''\colon L \to P$ *is trivial and* L *is an abelian Lie superalgebra.*

Proof. Clearly $\partial'' = \partial' \circ g \circ f = 0$ and $[l,l'] = {}^{\partial''(l)} l' = 0$, for all $l, l' \in L$. □

If (M,∂) and (N,∂') are two crossed P-modules, then the Lie superalgebras M and N act compatibly on each other via the action of P. Thus, we can construct the non-abelian tensor product of Lie superalgebras $M \otimes N$. Moreover, we have an action of P on $M \otimes N$ defined by ${}^p(m \otimes n) = {}^p m \otimes n + (-1)^{|p||m|} m \otimes {}^p n$, and straightforward computations show that $\eta\colon M \otimes N \to P$, $m \otimes n \mapsto [\partial(m), \partial'(n)]$, is a crossed P-module.

Proposition 5.2. *Let* (M,∂) *be a crossed* P*-module. There is a right exact functor* $(M \otimes -)\colon \mathtt{Cross}(\mathrm{P}) \to \mathtt{Cross}(\mathrm{P})$ *given, for any crossed* P*-module* (N,∂')*, by*

$$(M \otimes -)(N,\partial') = (M \otimes N, \eta).$$

Proof. It is an immediate consequence of Proposition 3.8 □

Definition 5.3. Let (M,∂) be a crossed P-module. We define the zero and first non-abelian homologies of P with coefficients in M by setting

$$\mathcal{H}_0(P,M) = \mathsf{Coker}\,\nu \qquad \text{and} \qquad \mathcal{H}_1(P,M) = \mathsf{Ker}\,\nu,$$

where $\nu\colon P \otimes M \to M$, $p \otimes m \mapsto {}^p m$, is the Lie superalgebra homomorphism as in Proposition 3.4.

If we consider the crossed P-module (P, id_P) we have that

$$\mathcal{H}_0(P,P) = \frac{P}{[P,P]} \cong \mathrm{H}_1(P).$$

In addition, if P is perfect, by Theorem 4.1 we have that $\mathcal{H}_1(P,P) \cong \mathrm{H}_2(P)$.

The zero and first non-abelian homologies generalize respectively the zero and first homologies of Lie superalgebras in the sense of the following proposition.

Proposition 5.4. *Let the ground ring $\mathbb{K}$ be a field. Let P be a Lie superalgebra and M a P-supermodule thought as a crossed P-module $(M, 0)$. Then there are isomorphisms of super vector spaces*

$$\mathcal{H}_0(P, M) \cong \mathrm{H}_0(P, M) \quad \textit{and} \quad \mathcal{H}_1(P, M) \cong \mathrm{H}_1(P, M).$$

Proof. This is a direct consequence of Proposition 3.6 and the isomorphisms (2) and (3). □

Proposition 5.5. *Given a short exact sequence in* $\mathtt{Cross}(\mathrm{P})$

$$0 \to (L, 0) \to (M, \partial) \to (N, \partial') \to 0$$

we have an exact sequence of supermodules

$$\mathcal{H}_1(P, L) \to \mathcal{H}_1(P, M) \to \mathcal{H}_1(P, N) \to \mathcal{H}_0(P, L) \to \mathcal{H}_0(P, M) \to \mathcal{H}_0(P, N) \to 0.$$

Proof. The proof is an immediate consequence of the snake lemma applied to the diagram obtained from Proposition 5.2

$$\begin{array}{ccccccccc} & & P \otimes L & \longrightarrow & P \otimes M & \longrightarrow & P \otimes N & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & L & \longrightarrow & M & \longrightarrow & N & \longrightarrow & 0. \end{array}$$

□

5.2. **Application to the cyclic homology of associative superalgebras.** Now we recall from [15] and [13] the definition of cyclic homology of associative superalgebras. Let A be an associative superalgebra and $(C'_*(A), d'_*)$ denote its Hochschild complex, that is $C'_n(A) = A^{\otimes_{\mathbb{K}}(n+1)}$ and the boundary map $d_n : C'_n(A) \to C'_{n-1}(A)$ is given by

$$d'_n(a_0 \otimes \cdots \otimes a_n) = \sum_{i=0}^{n-1} (-1)^i a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n + (-1)^{n+|a_n|(|a_0|+\cdots+|a_{n-1}|)} a_n a_0 \otimes \cdots \otimes a_{n-1}.$$

Now the cyclic group $\mathbb{Z}/(n+1)\mathbb{Z}$ acts on $A^{\otimes_{\mathbb{K}}(n+1)}$ via

$$t_n(a_0 \otimes \cdots \otimes a_n) = (-1)^{n+|a_n| \sum_{k<n} |a_k|} a_n \otimes a_0 \otimes \cdots \otimes a_{n-1},$$

where $t_n = 1 + (n+1)\mathbb{Z} \in \mathbb{Z}/(n+1)\mathbb{Z}$. For each $n \geq 0$, consider the quotient $C_n(A) = A^{\otimes_{\mathbb{K}}(n+1)} / \mathsf{Im}(1 - t_n)$ which is the module of coinvariants of $C'_n(A)$ under the $\mathbb{Z}/(n+1)\mathbb{Z}$-action. Then d'_n induces a well-defined map $d_n : C_n(A) \to C_{n-1}(A)$ and there is an induced chain complex $(C_*(A), d_*)$, which is called the Connes complex of A. Its homologies are, by definition, the *cyclic homologies of the associative superalgebra* A, denoted by $\mathrm{HC}_n(A)$, $n \geq 0$.

Easy calculations show that, given an associative superalgebra A, $\mathrm{HC}_1(A)$ is the kernel of the homomorphism of supermodules

$$(A \otimes_{\mathbb{K}} A)/\mathrm{I}(A) \to [A, A], \quad a \otimes b \mapsto ab - (-1)^{|a||b|} ba,$$

where $[A, A]$ is the graded submodule of A generated by the elements $ab - (-1)^{|a||b|} ba$ and $\mathrm{I}(A)$ is the graded submodule of the supermodule $A \otimes_{\mathbb{K}} A$ generated by the

elements

$$a \otimes b + (-1)^{|a||b|} b \otimes a,$$
$$ab \otimes c - a \otimes bc + (-1)^{|c|(|a|+|b|)} ca \otimes b,$$

for all homogeneous $a, b, c \in A$.

Now let us consider A as a Lie superalgebra (see Example 2.2(i)). Then there is a Lie superalgebra structure on $(A \otimes_{\mathbb{K}} A)/\mathrm{I}(A)$ given by

$$[a \otimes b, a' \otimes b'] = [a, b] \otimes [a', b']$$

for all $a, a', b, b' \in A$. We denote this Lie superalgebra by $\mathrm{V}(A)$. In fact, $\mathrm{V}(A)$ is the quotient of the non-abelian tensor product $A \otimes A$ by the graded ideal generated by the elements $x \otimes y + (-1)^{|x||y|} y \otimes x$ and $xy \otimes z - x \otimes yz + (-1)^{|z|(|x|+|y|)} zx \otimes y$, for all homogeneous $x, y, z \in A$.

Proposition 5.6. *Let A be a Lie superalgebra. Then the following assertions hold:*

(i) *There are compatible actions of the Lie superalgebras A and $\mathrm{V}(A)$ on each other.*
(ii) *The map $\mu\colon \mathrm{V}(A) \to A$ given by $x \otimes y \mapsto [x, y]$, together with the action of A on $\mathrm{V}(A)$, is a crossed module of Lie superalgebras.*
(iii) *The action of A on $\mathrm{V}(A)$ induces the trivial action of A on $\mathrm{HC}_1(A)$.*
(iv) *There is a short exact sequence in the category* `Cross`(A)

$$0 \to (\mathrm{HC}_1(A), 0) \to (\mathrm{V}(A), \mu) \to ([A, A], i) \to 0,$$

where $i\colon [A, A] \to A$ is the inclusion.

Proof.

(i) The action of A on $\mathrm{V}(A)$ is induced by the action of A on $A \otimes A$ given in Proposition 3.4(ii), that is

$$\begin{aligned} {}^a(x \otimes y) &= [a, x] \otimes y + (-1)^{|a||x|} x \otimes [a, y] \\ &= ax \otimes y + (-1)^{|a|(|x|+|y|)} x \otimes ya - (-1)^{|x||a|} x \otimes ay - (-1)^{|x||a|} xa \otimes y \\ &= a \otimes xy - (-1)^{|x||y|} a \otimes yx \\ &= a \otimes [x, y], \end{aligned}$$

whilst the action of $\mathrm{V}(A)$ on A is defined by

$${}^{x \otimes y}a = \big[[x, y], a\big]$$

for all homogeneous $a, x, y \in A$. Straightforward calculations show that these are indeed (compatible) actions of Lie superalgebras.

(ii) Since the crossed module of Lie superalgebras $A \otimes A \to A$, $x \otimes y \mapsto [x, y]$, given in Proposition 3.4, vanishes on the elements of the form $x \otimes y + (-1)^{|x||y|} y \otimes x$ and $xy \otimes z - x \otimes yz + (-1)^{|z|(|x|+|y|)} zx \otimes y$, then μ is well defined and obviously it is a crossed module of Lie superalgebras.

(iii) If $\sum_i \lambda_i (x_i \otimes y_i) \in \mathrm{HC}_1(A)$, i.e. $\sum_i \lambda_i [x_i, y_i] = 0$, then for all $a \in A$ we have

$${}^a\Big(\sum_i \lambda_i (x_i \otimes y_i)\Big) = \sum_i \lambda_i (a \otimes [x_i, y_i]) = a \otimes \sum_i \lambda_i [x_i, y_i] = 0.$$

(iv) This is an immediate consequence of the assertions above. □

By Proposition 5.5 we have the following exact sequence of supermodules

$$\begin{aligned}&\mathcal{H}_1\big(A,\mathrm{HC}_1(A)\big)\longrightarrow\mathcal{H}_1\big(A,\mathrm{V}(A)\big)\longrightarrow\mathcal{H}_1(A,[A,A])\\&\hookrightarrow\mathcal{H}_0\big(A,\mathrm{HC}_1(A)\big)\longrightarrow\mathcal{H}_0\big(A,\mathrm{V}(A)\big)\longrightarrow\mathcal{H}_0(A,[A,A])\longrightarrow 0.\end{aligned}\tag{6}$$

Below, we will calculate some of the terms of this exact sequence. At first, by analogy to the Dennis-Stein generators [6], we give a definition of the first Milnor cyclic homology for associative superalgebras.

Definition 5.7. Let A be an associative superalgebra. We define the first Milnor cyclic homology $\mathrm{HC}_1^M(A)$ of A to be the quotient of the supermodule $A\otimes_{\mathbb{K}}A$ by the graded ideal generated by the elements

$$\begin{aligned}&a\otimes b+(-1)^{|a||b|}b\otimes a,\\&ab\otimes c-a\otimes bc+(-1)^{|c|(|a|+|b|)}ca\otimes b,\\&a\otimes bc-(-1)^{|b||c|}a\otimes cb,\end{aligned}$$

for all homogeneous $a,b,c\in A$.

It is clear that if A is supercommutative, that is, $ab=(-1)^{|a||b|}ba$, for all homogeneous $a,b\in A$, then $\mathrm{HC}_1(A)\cong\mathrm{HC}_1^M(A)$.

Lemma 5.8. *We have the following equalities and isomorphisms*

(i) $\mathcal{H}_0\big(A,\mathrm{HC}_1(A)\big)=\mathrm{HC}_1(A)$,
(ii) $\mathcal{H}_1\big(A,\mathrm{HC}_1(A)\big)\cong A/[A,A]\otimes_{\mathbb{K}}\mathrm{HC}_1(A)$,
(iii) $\mathcal{H}_0(A,[A,A])=[A,A]/\big[A,[A,A]\big]$,
(iv) $\mathcal{H}_0\big(A,\mathrm{V}(A)\big)\cong\mathrm{HC}_1^M(A)$.

Proof.

(i) Since A acts trivially on $\mathrm{HC}_1(A)$, we have that $\mathsf{Coker}\big(A\otimes\mathrm{HC}_1(A)\to\mathrm{HC}_1(A)\big)=\mathrm{HC}_1(A)$.

(ii) Since $\mathrm{HC}_1(A)$ is abelian, by Proposition 3.5 we have that $\mathsf{Ker}\big(A\otimes\mathrm{HC}_1(A)\to\mathrm{HC}_1(A)\big)\cong A/[A,A]\otimes_{\mathbb{K}}\mathrm{HC}_1(A)$.

(iii) and (iv) are straightforward. □

It follows that the exact sequence (6) can be written as in the following theorem.

Theorem 5.9. *If A is a unital associative superalgebra. Then there is an exact sequence of supermodules*

$$\begin{aligned}&\frac{A}{[A,A]}\otimes_{\mathbb{K}}\mathrm{HC}_1(A)\longrightarrow\mathcal{H}_1\big(A,\mathrm{V}(A)\big)\longrightarrow\mathcal{H}_1(A,[A,A])\\&\hookrightarrow\mathrm{HC}_1(A)\longrightarrow\mathrm{HC}_1^M(A)\longrightarrow\frac{[A,A]}{\big[A,[A,A]\big]}\longrightarrow 0.\end{aligned}$$

Corollary 5.10. *If A is perfect as a Lie superalgebra, we have an exact sequence*

$$0\to\mathcal{H}_1\big(A,\mathrm{V}(A)\big)\to\mathrm{H}_2(A)\to\mathrm{HC}_1(A)\to 0,$$

where $\mathrm{H}_2(A)$ *is the usual second homology of the Lie superalgebra* A*. If in addition* $\mathrm{H}_2(A) = 0$*, then all terms of the exact sequence in the previous theorem are trivial.*

Proof. Since A is perfect we know that $\mathcal{H}_1(A, A) \cong \mathrm{H}_2(A)$, $A/[A, A] \otimes_{\mathbb{K}} \mathrm{HC}_1(A) = 0$ and the map $A \otimes \mathrm{V}(A) \to \mathrm{V}(A)$ is surjective. □

6. Non-abelian exterior product of Lie superalgebras

In this section we extend to Lie superalgebras the definition of the non-abelian exterior product of Lie algebras introduced in [8]. Then we use it to derive the Hopf formula for the second homology of a Lie superalgebra and to construct a six-term exact homology sequence of Lie superalgebras.

6.1. **Construction of the non-abelian exterior product.** Let P be a Lie superalgebra and (M, ∂) and (N, ∂') two crossed P-modules. We consider the actions of M and N on each other via P.

Lemma 6.1. *Let* $M \square N$ *be the graded submodule of* $M \otimes N$ *generated by the elements*

(a) $m \otimes n + (-1)^{|m'||n'|} m' \otimes n'$*, where* $\partial(m) = \partial'(n')$ *and* $\partial(m') = \partial'(n)$*,*
(b) $m_{\bar{0}} \otimes n_{\bar{0}}$*, where* $\partial(m_{\bar{0}}) = \partial'(n_{\bar{0}})$*,*

with $m, m' \in M_{\bar{0}} \cup M_{\bar{1}}$*,* $n, n' \in N_{\bar{0}} \cup N_{\bar{1}}$*,* $m_{\bar{0}} \in M_{\bar{0}}$ *and* $n_{\bar{0}} \in N_{\bar{0}}$*. Then,* $M \square N$ *is a graded ideal in the centre of* $M \otimes N$*.*

Proof. Given an element $m \otimes n + (-1)^{|m'||n'|} m' \otimes n'$ of the form (a), suppose that $|m'| = |n|$, then we have

$$\begin{aligned}
[x \otimes y, m \otimes n + (-1)^{|m'||n'|} m' \otimes n'] &= -(-1)^{|x||y|} ({}^{y}x) \otimes \left({}^{m}n + (-1)^{|m'||n'|} ({}^{m'}n') \right) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes \left({}^{\partial(m)}n + (-1)^{|m'||n'|} ({}^{\partial(m')}n') \right) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes \left({}^{\partial'(n')}n + (-1)^{|m'||n'|} ({}^{\partial'(n)}n') \right) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes \left([n', n] + (-1)^{|n||n'|} [n, n'] \right) \\
&= 0.
\end{aligned}$$

This is also true when $|m'| \neq |n|$. Indeed, if $|m'| \neq |n|$, since ∂, ∂' are even maps, the equality $\partial(m) = \partial'(n')$ holds if and only if $\partial(m) = 0 = \partial'(n')$. Now take an element $m_{\bar{0}} \otimes n_{\bar{0}}$ of the form (b). Then we have

$$\begin{aligned}
[x \otimes y, m_{\bar{0}} \otimes n_{\bar{0}}] &= -(-1)^{|x||y|} ({}^{y}x) \otimes ({}^{m_{\bar{0}}}n_{\bar{0}}) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes ({}^{\partial(m_{\bar{0}})}n_{\bar{0}}) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes ({}^{\partial(n_{\bar{0}})}n_{\bar{0}}) \\
&= -(-1)^{|x||y|} ({}^{y}x) \otimes [n_{\bar{0}}, n_{\bar{0}}] \\
&= 0,
\end{aligned}$$

for any $x \otimes y \in M \otimes N$. This completes the proof. □

Definition 6.2. Let P be a Lie superalgebra and (M, ∂) and (N, ∂') two crossed P-modules. The *non-abelian exterior product* $M \wedge N$ of the Lie superalgebras M and N is defined by

$$M \wedge N = \frac{M \otimes N}{M \square N}.$$

The equivalence class of $m \otimes n$ will be denoted by $m \wedge n$.

Note that if $M = M_{\bar{0}}$ and $N = N_{\bar{0}}$ then $M \wedge N$ coincides with the non-abelian exterior product of Lie algebras [8].

Reviewing Section 3, one can easily check that most of results on the non-abelian tensor product are fulfilled for the non-abelian exterior product. In particular, there are homomorphisms of Lie superalgebras $M \wedge N \to M$, $M \wedge N \to N$ and actions of M and N on $M \wedge N$, induced respectively by the homomorphisms and actions given in Proposition 3.4. It is also satisfied the isomorphism $M \wedge N \cong N \wedge M$. Further, given a short exact sequence of Lie superalgebras $0 \to K \to M \to P \to 0$, as an exterior analogue of the exact sequence (5), we get the following exact sequence of Lie superalgebras

$$K \wedge M \to M \wedge M \to P \wedge P \to 0. \tag{7}$$

Given a Lie superalgebra M, since $\mathsf{id}\colon M \to M$ is a crossed module, we can consider $M \wedge M$. It is the quotient of $M \otimes M$ by the following relations

$$\begin{aligned} m \wedge m' &= -(-1)^{|m||m'|} m' \wedge m, \\ m_{\bar{0}} \wedge m_{\bar{0}} &= 0, \end{aligned}$$

for all $m, m' \in M_{\bar{0}} \cup M_{\bar{1}}$ and $m_{\bar{0}} \in M_{\bar{0}}$. In the particular case when M is perfect, it is easy to see that $M \square M = 0$, so $M \wedge M \cong M \otimes M$ and in Theorem 4.1 we can replace $M \otimes M$ by $M \wedge M$.

6.2. **A six term exact homology sequence.** In [7], the non-abelian exterior product of Lie algebras is used to construct a six-term exact sequence of homology of Lie algebras. In this section we will extend these results to Lie superalgebras.

First of all, we prove an analogue of Miller's theorem [17] on free Lie superalgebras extending the similar result obtained in [7] for Lie algebras.

Proposition 6.3. *Let $F = \mathrm{F}(X)$ be the free Lie superalgebra on a graded set X. Then the homomorphism $F \wedge F \to F$, $x \wedge y \mapsto xy$ is injective.*

Proof. Let us prove that $[F, F] \cong F \wedge F$. Using the same notations as in Construction 2.9, we define a map $\phi\colon \mathrm{alg}(X) * \mathrm{alg}(X) \to F \wedge F$ by $\sum_i \lambda_i x_i y_i \mapsto \sum_i \lambda_i (x_i \wedge y_i)$, where $\mathrm{alg}(X) * \mathrm{alg}(X)$ is the free product of superalgebras. It is easy to see that ϕ is a $\mathbb{K}$-superalgebra homomorphism since $[x \wedge y, x' \wedge y'] = xy \wedge x'y'$. The ideal I is contained in $\mathrm{alg}(X) * \mathrm{alg}(X)$ and by using the defining relations of $F \wedge F$ it is not difficult to check that ϕ vanishes on I. So we have an induced map from $[F, F]$ to $F \wedge F$, which is inverse to the homomorphism $F \wedge F \to [F, F]$, $x \wedge y \mapsto xy$. □

Let P be a Lie superalgebra and take the quotient supermodule $(P \wedge_{\mathbb{K}} P)/\operatorname{Im} d_3$, where $d_3\colon \bigwedge^3_{\mathbb{K}}(P) \to \bigwedge^2_{\mathbb{K}}(P)$ is the boundary map in the homology complex $(C_*(P, \mathbb{K}), d_*)$. Here $\mathbb{K}$ is considered as a trivial P-module. We define a bracket in $(P \wedge_{\mathbb{K}} P)/\operatorname{Im} d_3$ by setting

$$[x \wedge y, x' \wedge y'] = [x, y] \wedge [x', y']$$

for all $x, y \in P$. As a particular case of the exterior analogue of Proposition 3.3 we have

Lemma 6.4. *There is an isomorphism of Lie superalgebras*

$$\frac{P \wedge_{\mathbb{K}} P}{\operatorname{Im} d_3} \approx P \wedge P.$$

Corollary 6.5.

(i) *For any Lie superalgebra P there is an isomorphism of supermodules*

$$H_2(P) \cong \mathsf{Ker}(P \wedge P \to P).$$

(ii) $H_2(F) = 0$ *if F is a free Lie superalgebra.*

(iii) *(Hopf Formula) Given a free presentation $0 \to R \to F \to P \to 0$ of a Lie superalgebra P, there is an isomorphism of supermodules*

$$H_2(P) \cong \frac{R \cap [F,F]}{[F,R]}.$$

Proof.

(i) This follows immediately from Lemma 6.4.

(ii) This is a consequence of (i) and Proposition 6.3.

(iii) Since $F \wedge F \cong [F,F]$, using the exact sequence (7), we have

$$P \wedge P \cong \frac{[F,F]}{[F,R]}.$$

Then Lemma 6.4 completes the proof. □

Theorem 6.6. *Let M be a graded ideal of a Lie superalgebra P. Then there is an exact sequence*

$$\mathsf{Ker}(P \wedge M \to P) \to \mathrm{H}_2(P) \to \mathrm{H}_2(P/M) \to \frac{M}{[P,M]} \to \mathrm{H}_1(P) \to \mathrm{H}_1(P/M) \to 0.$$

Proof. By using the exact sequence (7) we have the following commutative diagram of Lie superalgebras with exact rows

$$\begin{array}{ccccccccc} & & M \wedge P & \longrightarrow & P \wedge P & \longrightarrow & \dfrac{P}{M} \wedge \dfrac{P}{M} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M & \longrightarrow & P & \longrightarrow & \dfrac{P}{M} & \longrightarrow & 0. \end{array}$$

Since $\mathsf{Coker}(M \wedge P \cong P \wedge M \to M) \cong M/[P,M]$ and $\mathsf{Coker}(P \wedge P \to P) \cong P/[P,P] \cong \mathrm{H}_1(P)$, then the assertion follows by using snake lemma and Corollary 6.5(i). □

In particular, if P is a Lie algebra and M is an ideal of P, then this sequence coincides with the six-term exact sequence in the homology of Lie algebras obtained in [7].

Acknowledgements

The authors wish to thank the anonymous referee for his help in improving the presentation of this paper. The authors were supported by Ministerio de Economía y Competitividad (Spain), grant MTM2013-43687-P (European FEDER support included). The first and third authors were also supported by Xunta de Galicia, grant GRC2013-045 (European FEDER support included). The first author was also supported by FPU scholarship, Ministerio de Educación, Cultura y Deporte (Spain). The second author was also supported by Xunta de Galicia, grant EM2013/016 (European FEDER support included) and Shota Rustaveli National Science Foundation, grant DI/12/5-103/11.

References

[1] R. Brown, J.-L. Loday, Van Kampen theorems for diagrams of spaces, Topology 26 (3) (1987) 311–335, with an appendix by M. Zisman.

[2] C. Carmeli, L. Caston, R. Fioresi, Mathematical foundations of supersymmetry, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2011.

[3] J. M. Casas, M. Ladra, Perfect crossed modules in Lie algebras, Comm. Algebra 23 (5) (1995) 1625–1644.

[4] J. Castiglioni, X. García-Martínez, M. Ladra, Universal central extensions of Lie–Rinehart algebras, arXiv:1403.7159 (2014).

[5] H. Chen, J. Sun, Universal central extensions of $sl_{m|n}$ over $\mathbb{Z}/2\mathbb{Z}$-graded algebras, J. Pure Appl. Algebra 219 (2015) 4278–4294.

[6] R. K. Dennis, M. R. Stein, K_2 of discrete valuation rings, Advances in Math. 18 (2) (1975) 182–238.

[7] G. J. Ellis, Nonabelian exterior products of Lie algebras and an exact sequence in the homology of Lie algebras, J. Pure Appl. Algebra 46 (2-3) (1987) 111–115.

[8] G. J. Ellis, A nonabelian tensor product of Lie algebras, Glasgow Math. J. 33 (1) (1991) 101–120.

[9] X. García-Martínez, M. Ladra, Universal central extensions of $\mathfrak{sl}(m, n, a)$ of small rank over associative superalgebras, arXiv:1405.4035 (2014).

[10] D. Guin, Cohomologie des algèbres de Lie croisées et K-théorie de Milnor additive, Ann. Inst. Fourier (Grenoble) 45 (1) (1995) 93–118.

[11] P. J. Hilton, U. Stammbach, A course in homological algebra, vol. 4 of Graduate Texts in Mathematics, 2nd ed., Springer-Verlag, New York, 1997.

[12] N. Inassaridze, E. Khmaladze, M. Ladra, Non-abelian homology of Lie algebras, Glasg. Math. J. 46 (2) (2004) 417–429.

[13] K. Iohara, Y. Koga, Second homology of Lie superalgebras, Math. Nachr. 278 (9) (2005) 1041–1053.

[14] V. G. Kac, Lie superalgebras, Advances in Math. 26 (1) (1977) 8–96.

[15] C. Kassel, A Künneth formula for the cyclic cohomology of $\mathbf{Z}/2$-graded algebras, Math. Ann. 275 (4) (1986) 683–699.

[16] C. Kassel, J.-L. Loday, Extensions centrales d'algèbres de Lie, Ann. Inst. Fourier (Grenoble) 32 (4) (1982) 119–142.

[17] C. Miller, The second homology group of a group; relations among commutators, Proc. Amer. Math. Soc. 3 (1952) 588–595.

[18] I. M. Musson, Lie superalgebras and enveloping algebras, vol. 131 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2012.

[19] E. Neher, An introduction to universal central extensions of Lie superalgebras, in: Groups, rings, Lie and Hopf algebras (St. John's, NF, 2001), vol. 555 of Math. Appl., Kluwer Acad. Publ., Dordrecht, 2003, pp. 141–166.

[20] A. R. Salemkar, H. Tavallaee, H. Mohammadzadeh, B. Edalatzadeh, On the non-abelian tensor product of Lie algebras, Linear Multilinear Algebra 58 (3-4) (2010) 333–341.

[21] M. Scheunert, The theory of Lie superalgebras. An introduction, vol. 716 of Lecture Notes in Mathematics, Springer, Berlin, 1979.

[22] J. Tanaka, On homology and cohomology of Lie superalgebras with coefficients in their finite-dimensional representations, Proc. Japan Acad. Ser. A Math. Sci. 71 (3) (1995) 51–53.

[23] T. Zhang, Z. Liu, Omni-Lie superalgebras and Lie 2-superalgebras, Front. Math. China 9 (5) (2014) 1195–1210.

Xabier García-Martínez: Department of Algebra, IMAT, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain

E-mail address: xabier.garcia@usc.es

Emzar Khmaladze: A. Razmadze Mathematical Institute of Tbilisi State University, Tamarashvili Str. 6, 0177 Tbilisi, Georgia

E-mail address: e.khmal@gmail.com

Manuel Ladra: Department of Algebra, IMAT, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain
E-mail address: manuel.ladra@usc.es